BMJ Publishing Group Limited (BMJ) disclaims all liability and responsibility arising from any reliance

Supplemental material   placed on this supplemental material which has been supplied by the author(s)   *Reg Anesth Pain Med*

@RAPMOnline

POORER OUTCOMES

For Hip Arthroplasty in High vs Low Safety Net Burden Hospitals

DOI: 10.1136/rapm-2018-100305

Safety Net Hospitals

"Hospitals with a high proportion of cases billed to Medicaid or without insurance, [who] deliver a significant portion of their care to vulnerable populations"

California
Florida
Maryland
New York
Kentucky

State Inpatient Databases

2007 - 2014

500,000+ Patients

Burden of Medicaid or Uninsured (%)

High	Medium	Low
> 30.5%	17-30.5%	<17%

High Safety Net Hospitals	VS	Low Safety Net Hospitals

↑	38%	worse	Mortality
↑	11%	worse	Postop Complications
↑	15%	worse	CV Complications
↑	24%	worse	Infectious Complications
↑	6%	worse	Length of Stay

High Safety Net Hospitals

2.5 TIMES

More likely to have

General Anesthesia

instead of

Regional Anesthesia

Click Here for Article

Regional Anesthesia & Pain Medicine

@RAPMOnline