

Supplement Table 2. Baseline characteristics of patients who underwent outpatient total knee or hip arthroplasty by same day discharge status.

Characteristic	Same Day Discharge Status			p-value
	Overall	Failed same day discharge	Same day discharge	
	N = 11523	N = 2838	N = 8685	
Mean Age in Years (SD)	68.0 (8.9)	69.7 (9.5)	67.5 (13.8)	<0.01 ^a
Sex, n (%)				
Male	4577 (39.7)	885 (31.2)	3692 (42.5)	<0.01
Female	6946 (60.3)	1953 (68.8)	4993 (57.5)	
Race/Ethnicity, n (%)				
Non-Hispanic White	7844 (68.1)	1801 (63.5)	6043 (69.6)	<0.01
Asian	931 (8.1)	275 (9.7)	656 (7.6)	
Black	720 (6.3)	222 (7.8)	498 (5.7)	
Hispanic	1515 (13.2)	404 (14.2)	1111 (12.8)	
Other	513 (4.5)	136 (4.8)	377 (4.3)	
BMI^b, n (%)				
Normal	1727 (15.0)	458 (16.1)	1269 (14.6)	0.09
Overweight	3967 (34.4)	943 (33.2)	3024 (34.8)	
Obese	5828 (50.6)	1437 (50.6)	4391 (50.6)	
ASA Class, n (%)				
I	172 (1.5)	28 (1.0)	144 (1.7)	<0.01
II	6874 (59.7)	1462 (51.5)	5412 (62.3)	
III	4396 (38.2)	1300 (45.8)	3096 (35.7)	
IV	81 (0.7)	48 (1.7)	33 (0.4)	
Surgery Type, n (%)				
Knee	9866 (85.6)	2398 (84.5)	7468 (86.0)	0.04
Hip	1657 (14.4)	440 (15.5)	1217 (14.0)	
Comorbidity Score^c, n (%)				
0	4412 (38.3)	894 (31.5)	3518 (40.5)	<0.01
1-2	4854 (42.1)	1230 (43.3)	3624 (41.7)	
3+	2257 (19.6)	714 (25.2)	1543 (17.8)	
Laterality, n (%)				
Left	5470 (47.5)	1320 (46.5)	4150 (47.8)	0.24
Right	6053 (52.5)	1518 (53.5)	4535 (52.2)	
Mean Hemoglobin Before Surgery within 3 months of surgery date^d (SD)	13.6 (1.4)	13.2 (1.5)	13.8 (1.3)	<0.01 ^a

Abbreviations: SD, standard deviation; BMI, body mass index; ASA, American Society of Anesthesiologists.

^ap-value calculated with Student t-test^b0.008% missing BMI (One patient)^cCharlson comorbidity index used as comorbidity score^d10.5% missing last hemoglobin before surgery (1214 patients)